

Aqua Neat[®]

Aquatic Herbicide

FOR USE ON EMERGED AQUATIC WEEDS AND BRUSH IN AQUATIC SITES. FOR USE IN FORESTRY (INCLUDING WEED CONTROL IN CHRISTMAS TREE PLANTATIONS), RIGHTS-OF-WAY, HABITAT RESTORATION AREAS, NON-CROP AND OTHER LISTED APPLICATION SITES.

ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl)glycine,
in the form of its isopropylamine salt*

53.8%

OTHER INGREDIENTS:

46.2%

TOTAL:

100.0%

*Contains 648 grams per litre or 5.4 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 480 grams per litre or 4 pounds per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND
ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or
Exposure, Call CHEMTREC
(800) 424-9300

For Medical Emergencies Only,
Call (877) 325-1840

EPA Reg. No. 228-365

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCION**

Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

FIRST AID	
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS
<p>Users Should:</p> <ul style="list-style-type: none">• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For aquatic uses, do not contaminate water when disposing of equipment washwaters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

In case of, SPILL OR LEAK, soak up and remove to a landfill. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE) and Restricted-Entry Interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the Restricted-Entry Interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, shoes plus socks, and chemical-resistant gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber) > 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

PRODUCT INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL OR CURRENT SUPPLEMENTAL LABELING ISSUED BY MANUFACTURER.

This product, a water-soluble liquid, mixes readily with water and nonionic surfactant to be applied as a foliar spray after dilution and thoroughly mixing with water in accordance with label instructions for the control or destruction of many herbaceous and woody plants. Always use the higher rate of this product per acre within the specified range when vegetation is heavy or dense. When treating dense multi-canopied sites or woody vegetation or difficult-to-control herbaceous or woody plants.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial brush species may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow the activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise directed on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label.

Unemerged plants arising from unattached underground rhizomes or root stocks of perennials or brush will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds or brush is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the specified range when vegetation is heavy or dense.

Do not treat weeds or brush under poor growing conditions such as drought stress, disease or insect damage, as reduced control may result. Reduced results may also occur when treating weeds or brush heavily covered with dust.

Reduced control may result when applications are made to any weed or brush species that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the product off the foliage and a repeat treatment may be required.

Mixing this product with herbicides or other materials not instructed in this label may result in reduced performance. However, unless otherwise prohibited on this label or the label of an intended tank mix product may be applied in combination with any herbicide registered for the same site, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. **TO THE FULLEST EXTENT PERMITTED BY LAW, BUYER AND ALL USERS ARE RESPONSIBLE FOR ALL LOSS OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF MIXTURES OF THIS PRODUCT OR OTHER MATERIALS THAT ARE NOT EXPRESSLY SPECIFIED IN THIS LABEL.**

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

When this product comes in contact with soil (on the soil surface or as suspended soil or sediment in water) it is bound to soil particles. Under labeled use situations, once this product is bound to soil particles, it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treatment area or if the soil is transported off-site. Under labeled use conditions, the strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water. The affinity between this product and soil particles remains until this product is degraded, which is primarily a biological degradation process carried out under both aerobic and anaerobic conditions by soil micro flora.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Read **“WARRANTY DISCLAIMER”** and **“LIMITATION OF LIABILITY”** before buying or using. If items are not acceptable, return at once unopened. Buyer and all users are responsible for all loss or damage in connection with the use of handling of mixtures of this product or other materials that are not expressly specified in this label.

For more product information, call toll-free 1-800-345-3330.

ATTENTION

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

MIXING AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. HAND-GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance. Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted SLOWLY through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Where nonionic surfactant is recommended, add this to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. To prevent or minimize foam, avoid the use of mechanical agitators, place the filling hose below the surface of the spray solution, terminate by-pass and return lines at the bottom of the tank and if needed use an approved anti-foam or defoaming agent.

Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Use a nonionic surfactant labeled for use with herbicides. The surfactant must contain 50 percent or more active ingredient.

Always read and follow the manufacturer's surfactant label instructions for best results.

These surfactants should not be used in excess of 1 quart per acre when making broadcast applications.

Colorants or marking dyes approved for use with herbicides may be added to spray mixtures of this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's label instructions.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water and dispose of rinsate according to labeled use or disposal instructions.

Carefully observe all cautionary statements and other information appearing in the surfactant label.

APPLICATION EQUIPMENT AND TECHNIQUES

This product may be applied with the following application equipment:

Aerial - Fixed Wing and Helicopter

Broadcast Spray

Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Hand-Held and High-Volume Spray Equipment* - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment - Recirculating sprayers and wiper applicators. See the appropriate part of this section for specific instructions and rates of application.

APPLICATION INFORMATION

Observe the following directions to minimize off-site movement during aerial application of this herbicide. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

AERIAL EQUIPMENT

Use the specified rates of this product and surfactant in 3 to 20 gallons of water per acre as a broadcast spray, unless otherwise specified. See the "WEEDS CONTROLLED" section of this label for specific rates. Unless otherwise specified, do not exceed 1.5 pints per acre. Aerial applications of this product may only be made as specified in this label.

AVOID DRIFT - DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure above the manufacturer's instructions.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing in the additive label.

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

For use of this product by air in California see additional instructions in "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" Section.

FOR AERIAL APPLICATION IN CALIFORNIA ONLY

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Written Directions

A written direction **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written direction **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this herbicide is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at night

Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

Aquatic and Other Noncrop Sites

When applied as directed and under the conditions described in the "Weeds Controlled" section of the label booklet for this product, this herbicide will control or partially control the labeled weeds growing in the following industrial, recreational and public areas, or other similar sites.

Aquatic Sites-including all bodies of fresh and brackish water which may be flowing, nonflowing or transient. This includes lakes, rivers, streams, ponds, seeps, irrigation and drainage ditches, canals, reservoirs, estuaries and similar sites.

If aquatic sites are present in the noncrop areas and are part of the intended treatment, read and observe the following directions: There is no limit on the use of treated water for irrigation, recreation or domestic purposes.

Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.

NOTE: Do not apply this product within 1/2 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as a lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made **ONLY** in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after application.

This product does not control plants which are completely submerged or have a majority of their foliage underwater.

AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

FOR AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

From February 15 through March 31 only. For aerial application outside of these dates, refer to the "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" section printed above.

APPLICABLE AREA

This supplement only applies to the area contained inside the following boundaries within Fresno County, California only.

- North: Fresno County line
- South: Fresno County line
- East: State Highway 99
- West: Fresno County line

BOOM EQUIPMENT

For control of weed or brush species listed in this label using conventional boom equipment - Use the specified rates of this product and surfactant in 3 to 30 gallons of water per acre as a broadcast spray, unless otherwise specified. See the "WEEDS CONTROLLED" section of this label for specific rates. As density of vegetation increases, spray volume should be increased within the specified range to ensure complete coverage. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

HAND-HELD AND HIGH-VOLUME EQUIPMENT

Use Coarse Sprays Only

For control of weeds listed in this label using knapsack sprayers or high-volume spraying equipment utilizing handguns or other suitable nozzle arrangements - Prepare a 0.75 to 2 percent solution of this product in water, add a nonionic surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section in this label.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

This product may be used as a 5 to 8 percent solution plus 0.5 to 1 fluid ounce non-ionic surfactant per gallon spray solution for low-volume directed sprays for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zig-zag motion. Ensure that at least 50 percent of the leaves are contacted by the spray solution. For flat fan and cone nozzles and with hand-directed mist blowers, mist the application over the foliage of the targeted vegetation. Small, open-branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

For use in knapsack sprayers, it is suggested that the specified amount of this product be mixed with water in a large container. Fill sprayer with the mixed solution and add the correct amount of surfactant.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

SPRAY SOLUTION

DESIRED VOLUME	AMOUNT OF PRODUCT					
	0.75%	1.0%	1.25%	1.5%	5.0%	8.0%
1 Gallon	1.0 fl. oz.	1.33 fl. oz.	1.66 fl. oz.	2.0 fl. oz.	6.0 fl. oz.	10.25 fl. oz.
25 Gallons	1.5 pts.	1.0 qt.	1.25 qts.	1.5 qts.	5.0 qts.	2.0 gals.
100 Gallons	3.0 qts.	1.0 gal.	1.25 gals.	1.5 gals.	5.0 gals.	8.0 gals.

2 Tablespoons = 1 fluid ounce

SELECTIVE EQUIPMENT

For terrestrial application, this product may be applied through a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any non-crop site specified on this label.

- A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.
- A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

AVOID CONTACT WITH DESIRABLE VEGETATION.

This section summarizes the general weed control spectrum and rates of application for this herbicide. Additional information specific to individual use patterns is detailed in following sections.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees. Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the [Aerial Drift Reduction Advisory Information](#).

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions sections of this label).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Higher pressure reduces droplet size and does not improve canopy protection.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released backwards, parallel to the air stream produces larger droplets than other orientations. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.
- **Boom Length** - For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application Height** - Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

WEEDS CONTROLLED

ANNUAL WEEDS

Apply to actively growing annual grasses and broadleaf weeds.

Allow at least 3 days after application before disturbing treated vegetation. After this period the weeds may be mowed, tilled or burned. See "DIRECTIONS FOR USE", "PRODUCT INFORMATION" and "MIXING AND APPLICATION INSTRUCTIONS" for labeled uses and specific application instructions.

Broadcast Application - Use 1-1/2 pints of this product per acre plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution, if weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 2-1/2 pints of this product per acre plus 2 or more quarts of an approved nonionic surfactant per 100 gallons of spray solution.

Hand-Held, High-Volume Application - Use a 3/4 percent solution of this product in water plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution and apply to foliage of vegetation to be controlled.

When applied as directed under the conditions described in this label, this product plus nonionic surfactant WILL CONTROL the following ANNUAL WEEDS:

Balsamapple**

Momordica charantia

Barley

Hordeum vulgare

Barnyardgrass

Echinochloa crus-galli

Bassia, fivehook

Bassia hyssopifolia

Bluegrass, annual

Poa annua

Bluegrass, bulbous

Poa bulbosa

Brome*

Bromus spp.

Buttercup

Ranunculus spp.

Cheat

Bromus secalinus

Chickweed, mouseear

Cerastium vulgatum

Cocklebur

Xanthium strumarium

Corn, volunteer

Zea mays

Crabgrass

Digitaria spp.

Dwarf dandelion

Krigia cespitosa

False dandelion

Krigia cespitosa

Falseflax, smallseed

Camelina microcarpa

Fiddleneck*

Amsinckia spp.

Flax leaf fleabane*

Conyza bonariensis

Fleabane

Erigeron spp.

Foxtail

Setaria spp.

Foxtail, Carolina

Alopecurus carolinianus

Groundsel, common

Senecio vulgaris

Horseweed/Marestail

Conyza canadensis

Kochia*

Kochia scoparia

Lambsquarters, common

Chenopodium album

Lettuce, prickly*

Lactuca serriola

Morningglory

Ipomoea spp.

Mustard, blue

Chorispora tenella

Mustard, tansy

Descurainia pinnata

Mustard, tumble

Sisymbrium altissimum

Mustard, wild

Sinapis arvensis

Oats, wild

Avena fatua

Panicum*

Panicum spp.

(continued)

Pennycress, field
Thlaspi arvense
Pigweed, redroot
Amaranthus retroflexus
Pigweed, smooth
Amaranthus hybridus
Ragweed, common*
Ambrosia artemisiifolia
Ragweed, giant*
Ambrosia trifida
Rocket, London
Sisymbrium irio
Rye
Secale cereale
Ryegrass, Italian*
Lolium multiflorum

Sandbur, field
Cenchrus spp.
Shattercane
Sorghum bicolor
Shepherd's-purse
Capsella bursa-pastoris
Signalgrass, broadleaf
Brachiaria platyphylla
Smartweed, Pennsylvania
Polygonum pennsylvanicum
Sowthistle, annual*
Sonchus oleraceus
Spanishneedles*
Bidens bipinnata
Spurry, umbrella
Holosteum umbellatum

Stinkgrass
Eragrostis ciliaris
Sunflower*
Helianthus annuus
Thistle, Russian
Salsola kali
Velvetleaf*
Abutilon theophrasti
Wheat
Triticum aestivum
Witchgrass
Panicum capillare

*Apply 3 pints of this product per acre.
**Apply with hand-held equipment only.

Annual weeds will generally continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating weeds.

PERENNIAL WEEDS

Apply this product as follows to control or destroy most vigorously growing perennial weeds. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

See individual control instructions for specific weeds following the table. For other perennials listed on this label, apply 4-1/2 to 7-1/2 pints of product per acre as a broadcast spray or as a 3/4 to 1-1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached early head or early bud stage of growth.

Add 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution to the rates of this product given in this list. See the "PRODUCT INFORMATION", "DIRECTIONS FOR USE" and "MIXING AND APPLICATION" sections in this label for specific uses and application instructions.

NOTE: If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

When applied as specified under the conditions described, this product plus surfactant WILL CONTROL the following PERENNIAL WEEDS:

Alfalfa
Medicago sativa
Alligatorweed*
Alternanthera philoxeroides
Anise/Fennel
Foeniculum vulgare
Artichoke, Jerusalem
Helianthus tuberosus
Bahiagrass
Paspalum notatum
Bermudagrass
Cynodon dactylon
Bindweed, field
Convolvulus arvensis
Bluegrass, Kentucky
Poa pratensis
Blueweed, Texas
Helianthus ciliaris
Brackenfern
Pteridium spp.
Bromegrass, smooth
Bromus inermis

Canarygrass, reed
Phalaris arundinacea
Cattail
Typha spp.
Clover, red
Trifolium pratense
Clover, white
Trifolium repens
Cogongrass
Imperata cylindrica
Cordgrass
Spartina spp.
Cutgrass, giant*
Zizaniopsis miliacea
Dallisgrass
Paspalum dilatatum
Dandelion
Taraxacum officinale
Dock, curly
Rumex crispus
Dogbane, hemp
Apocynum cannabinum

Fescue
Festuca spp.
Fescue, tall
Festuca arundinacea
Guineagrass
Panicum maximum
Hemlock, poison
Conium maculatum
Horsenettle
Solanum carolinense
Horseradish
Armoracia rusticana
Ice Plant
Mesembryanthemum crystallinum
Johnsongrass
Sorghum halepense
Kikuyugrass
Pennisetum clandestinum
Knapweed
Centaurea repens
Lantana
Lantana camara

(continued)

Lespedeza: common, services

Lespedeza striata
Lespedeza cuneata

Loosestrife, purple

Lythrum salicaria

Lotus, American

Nelumbo lutea

Maidencane

Panicum hematomon

Milkweed

Asclepias spp.

Muhly, wirestem

Muhlenbergia frondosa

Mullein, common

Verbascum thapsus

Napiergrass

Pennisetum purpureum

Nightshade, silverleaf

Solanum elaeagnifolium

Nutsedge: purple, yellow

Cyperus rotundus
Cyperus esculentus

Orchardgrass

Dactylis glomerata

Pampas grass

Cortaderia jubata

Paragrass

Brachiaria mutica

Phragmites**

Phragmites spp.

Quackgrass

Agropyron repens

Reed, giant

Arundo donax

Ryegrass, perennial

Lolium perenne

Smartweed, swamp

Polygonum coccineum

Spatterdock

Nuphar luteum

Starthistle, yellow

Centaurea solstitialis

Sweet potato, wild*

Ipomoea pandurata

Thistle, artichoke

Cynara cardunculus

Thistle, Canada

Cirsium arvense

Timothy

Phleum pratense

Torpedograss*

Panicum repens

Tules, common

Scirpus acutus

Vaseygrass

Paspalum urvillei

Velvetgrass

Holcus spp.

Waterhyacinth

Eichornia crassipes

Waterlettuce

Pistia stratiotes

Waterprimrose

Ludwigia spp.

Wheatgrass, western

Agropyron smithii

*Partial control.

**Partial control in southeastern states. See specific instructions below.

Alligatorweed - Apply 6 pints of this product per acre as a broadcast spray or as a 1-1/4 percent solution with hand-held equipment to provide partial control of alligatorweed. Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control.

Bermudagrass - Apply 7-1/2 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and when seedheads appear.

Bindweed, field/Silverleaf Nightshade/Texas Blueweed - Apply 6 to 7-1/2 pints of this product per acre as a broadcast spray west of the Mississippi River and 4-1/2 to 6 pints of this product per acre east of the Mississippi River. With hand-held equipment, use a 1-1/2 percent solution. Apply when target plants are actively growing and are at or beyond full bloom. For silverleaf nightshade, best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall.

Brackenfern - Apply 4-1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Cattail - Apply 4-1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the early-to-full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.

Cogongrass - Apply 4-1/2 to 7-1/2 pints of this product per acre as a broadcast spray. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Cordgrass - Apply 4-1/2 to 7-1/2 pints of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment. Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. The presence of debris and silt on the cordgrass plants will reduce performance. It may be necessary to wash targeted plants prior to application to improve uptake of this product into the plant.

Cutgrass, giant - Apply 6 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain such control, especially where vegetation is partially submerged in water. Allow for substantial regrowth to the 7- to 10-leaf stage prior to retreatment.

Dogbane, hemp/Knapweed/Horseradish - Apply 6 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth. For best results, apply in late summer or fall.

Fescue, tall - Apply 4-1/2 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained.

Guineagrass - Apply 4-1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and when most have reached at least the 7-leaf stage of growth.

Johnsongrass/Bluegrass, Kentucky/Bromegrass, smooth/Canarygrass, reed/Orchardgrass/Ryegrass, perennial/Timothy/Wheatgrass, western - Apply 3 to 4-1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained. In the fall, apply before plants have turned brown.

Lantana - Apply this product as a 3/4 to 1 percent solution with hand-held equipment. Apply to actively growing Lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Loosestrife, purple - Apply 4 pints of this product per acre as a broadcast spray or as a 1 to 1-1/2 percent solution using hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost.

Lotus, American - Apply 4 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatment may be necessary to control regrowth from underground parts and seeds.

Maidencane/Paragrass - Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Repeat treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the 7- to 10-leaf stage prior to retreatment.

Milkweed, common - Apply 4-1/2 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth.

Nutsedge: purple, yellow - Apply 4-1/2 pints of this product per acre as a broadcast spray, or as a 3/4 percent solution with hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.

Pampasgrass - Apply a 1-1/2 percent solution of this product with hand-held equipment when plants are actively growing.

Phragmites - For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 7-1/2 pints per acre as a broadcast spray or apply a 1-1/2 percent solution with hand-held equipment. In other areas of the U.S., apply 4 to 6 pints per acre as a broadcast spray or apply a 3/4 percent solution with hand-held equipment for partial control. For best results, treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation, which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

Quackgrass/Kikuyugrass/Muhly, wirestem - Apply 3 to 4-1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches in height (3- to 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Reed, giant/ice plant - For control of giant reed and ice plant, apply a 1-1/2 percent solution of this product with hand-held equipment when plants are actively growing. For giant reed, best results are obtained when applications are made in late summer to fall.

Spatterdock - Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.

Sweet potato, wild - Apply this product as a 1-1/2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment.

Thistle: Canada, artichoke - Apply 3 to 4-1/2 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment for Canada thistle. To control artichoke thistle, apply a 2 percent solution as a spray to wet application. Apply when target plants are actively growing and are at or beyond the bud stage of growth.

Torpedograss - Apply 6 to 7-1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1-1/2 percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or a floating mat condition. Repeat treatments will be required to maintain such control.

Tules, common - Apply this product as a 1-1/2 percent solution with hand-held equipment. Apply to actively growing plants at or beyond the seedhead stage of growth. After application, visual symptoms will be slow to appear and may not occur for 3 or more weeks.

Waterhyacinth - Apply 5 to 6 pints of this product per acre as a broadcast spray or apply a 3/4 to 1 percent solution with hand-held equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are desired.

Waterlettuce - For control, apply a 3/4 to 1 percent solution using hand-held equipment to actively growing plants. Use higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring applications may require retreatment.

Waterprimrose - Apply this product as a 3/4 percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control.

Other perennials listed on this label - Apply 4-1/2 to 7-1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1-1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached early head or early bud stage of growth.

WOODY BRUSH AND TREES

See individual control instructions for specific woody brush and trees to be controlled in the following table. For partial control of other woody brush and trees listed in the table, apply 1.5 to 7.5 quarts of this product per acre as a broadcast spray or as a 0.75 to 10 percent solution with hand-held equipment.

Apply the specified rate of this product plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution when plants are actively growing and, unless otherwise directed, after full-leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late Summer or Fall after fruit formation.

Applied as a 5 to 8 percent solution as a directed application as described in the "HAND-HELD AND HIGH-VOLUME EQUIPMENT" section, this product will control or partially control all species listed in this section of the label. Use the higher rate of application for dense stands and larger woody brush and trees.

In arid areas, best results are obtained when application is made in the Spring or early Summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with Fall treatment.

Allow 7 or more days after application before mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if Fall treatments are made following a frost.

Application Rates¹

METHOD OF APPLICATION	APPLICATION RATE	SPRAY VOLUME (Gallons/Acre)
Broadcast Aerial Ground	1.5 to 7.5 qts./ acre 1.5 to 7.5 qts./ acre	5 to 30 10 to 60
Spray-to-Wet Handgun, Backpack, Mistblower	0.75% to 2.0% by volume	Spray-to-Wet
Low Volume Directed Spray² Handgun, Backpack, Mistblower	5.0% to 10.0% by volume	Partial Coverage

¹ Where repeat applications are necessary do not exceed 8.0 quarts per acre per year.

² For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. For best results, coverage of the top one-half of the plant is important.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stage of growth.

When applied as specified under the conditions described, this product plus surfactant CONTROLS or PARTIALLY CONTROLS the following woody brush plants and trees:

Alder

Alnus spp.

Ash*

Fraxinus spp.

Aspen, quaking

Populus tremuloides

Bearclover, Bearmat

Chamaebatia foliolosa

Birch

Betula spp.

Blackberry

Rubus spp.

Broom:

Cytisus monspessulanus

Scotch

Cytisus scoparius

Buckwheat, California*

Eriogonum fasciculatum

Cascara*

Rhamnus purshiana

Catsclaw*

Acacia greggi

Ceanothus

Ceanothus spp.

Chamise

Adenostoma fasciculatum

Cherry:

Bitter

Prunus emarginata

Black

Prunus serotina

Pin

Prunus pensylvanica

Coyote brush

Baccharis consanguinea

Creeper, Virginia*

Parthenocissus quinquefolia

Dewberry

Rubus trivialis

Dogwood

Cornus spp.

Elderberry

Sambucus spp.

Elm*

Ulmus spp.

Eucalyptus, bluegum

Eucalyptus globules

Hasardia*

Haplopappus squamosus

Hawthorn

Crataegus spp.

Hazel

Corylus spp.

Hickory

Carya spp.

Holly, Florida; Brazilian Peppertree

Schinus terebinthifolia

Honeysuckle

Lonicera spp.

Hornbeam, American

Carpinus caroliniana

Kudzu
Pueraria lobata

Locust, black*
Robinia pseudoacacia

Manzanita
Arctostaphylos spp.

Maple:
Red**
Acer rubrum

Sugar
Acer saccharum

Vine*
Acer circinatum

Monkey Flower*
Mimulus guttatus

Oak:
Black*
Quercus velutina

Northern pine
Quercus palustris

Post
Quercus stellata

Red
Quercus rubra

Southern red
Quercus falcata

White*
Quercus alba

Persimmon*
Diospyros spp.

Poison Ivy
Rhus radicans

Poison Oak
Rhus toxicodendron

Poplar, yellow*
Liriodendron tulipifera

Prunus
Prunus spp.

Raspberry
Rubus spp.

Redbud, eastern
Cercis canadensis

Rose, multiflora
Rosa multiflora

Russian-olive
Elaeagnus angustifolia

Sage: black, white
Salvia spp.

Sagebrush, California
Artemisia californica

Salmonberry
Rubus spectabilis

Salt cedar*
Tamarix spp.

Saltbush, Sea myrtle
Baccharis halimifolia

Sassafras
Sassafras albidum

Sourwood*
Oxydendrum arboreum

Sumac:
Poison*
Rhus vernix

Smooth*
Rhus glabra

Winged*
Phus copallina

Sweet gum
Liquidambar styraciflua

Swordfern*
Polystichum munitum

Tallowtree, Chinese
Sapium sebiferum

Thimbleberry
Rubus parviflorus

Tobacco, tree*
Nicotiana glauca

Trumpet creeper
Campsis radicans

Waxmyrtle, southern*
Myrica cerifera

Willow
Salix spp.

*Partial control

**See below for control or partial control instruction.

See the "DIRECTIONS FOR USE" and "MIXING AND APPLICATION INSTRUCTIONS" sections in this label for labeled use and specific application instructions.

Apply the product as follows to control or partially control the following woody brush and trees.

Alder/Blackberry/Dewberry/Honeysuckle/Oak, Post/Raspberry - For control, apply 4-1/2 to 6 pints per acre as a broadcast spray or as a 3/4 to 1-1/4 percent solution with hand-held equipment.

Aspen, Quaking/Hawthorn/Trumpet creeper - For control, apply 3 to 4-1/4 pints of this product per acre as a broadcast spray or as a 3/4 to 1-1/4 percent solution with hand-held equipment.

Birch/Elderberry/Hazel/Salmonberry/Thimbleberry - For control, apply 3 pints per acre of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment.

Broom: French, Scotch - For control, apply a 1-1/4 to 1-1/2 percent solution with hand-held equipment.

Buckwheat, California/Hasardia/Monkey Flower/Tobacco, Tree - For partial control of these species apply a 3/4 to 1-1/2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Catsclaw - For partial control, apply a 1-1/4 to 1-1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Cherry: Bitter, Black, Pin/Oak, Southern Red/Sweet Gum/Prunus - For control, apply 3 to 7-1/2 pints of this product per acre as a broadcast spray or as a 1 to 1-1/2 percent solution with hand-held equipment.

Coyote brush - For control, apply a 1-1/4 to 1-1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Dogwood/Hickory/Salt cedar - For partial control, apply a 1 to 2 percent solution of this product with hand-held equipment or 6 to 7-1/2 pints per acre as a broadcast spray.

Eucalyptus, bluegum - For control of eucalyptus resprouts, apply a 1-1/2 percent solution of this product with hand-held equipment when resprouts are 6- to 12-feet tall. Ensure complete coverage. Apply when plants are actively growing. Avoid application to drought-stressed plants.

Holly, Florida/Waxmyrtle, southern - For partial control, apply this product as a 1-1/2 percent solution with hand-held equipment.

Kudzu - For control, apply 6 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

Maple, Red - For control, apply as a 3/4 to 1-1/4 percent solution with hand-held equipment when leaves are fully developed. For partial control, apply 2 to 7-1/2 pints of this product per acre as a broadcast spray.

Maple, Sugar/Oak: Northern Pine, Red - For control, apply as a 3/4 to 1-1/4 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Poison Ivy/Poison Oak - For control, apply 6 to 7-1/2 pints of this product per acre as a broadcast spray or as a 1-1/2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Rose, multiflora - For control, apply 3 pints of this product per acre as a broadcast spray or as a 3/4 to 1-1/2 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

Sage, black/Sagebrush, California/Chamise/Tallowtree, Chinese - For control of these species, apply a 3/4 percent solution with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Saltbush, Sea myrtle - For control, apply this product as a 1 percent solution with hand-held equipment.

Willow - For control, apply 4-1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment.

Other woody brush and trees listed in this label - For partial control, apply 3 to 7-1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1-1/2 percent solution with hand-held equipment.

NON-CROP USES

See "PRODUCT INFORMATION" and "MIXING AND APPLICATION INSTRUCTIONS" sections of this label for essential product performance information and the following "NON-CROP" sections for specific uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OR SPRAY WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE TURFGRASSES, TREES, SHRUBS OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Where repeat applications are necessary, do not exceed 8 quarts of this product per acre per year.

This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

INDUSTRIAL, RECREATIONAL AND PUBLIC AREAS

When applied as directed for "NON-CROP USES", under conditions described, this product may be used to control the listed weeds.

Non-Crop Sites - This product may be used to control the listed weeds in terrestrial noncrop sites and/or in aquatic sites within these areas:

airfields; airports; alleys, lanes, trails & access roads; around commercial or industrial structures or outbuildings; around farm and ranch structures and outbuildings; around ornamental gardens; around ornamental trees & shrubs; bare ground; beaches; campgrounds; construction sites; ditch banks; drive-in theaters; driveways & ramps; dry ditches & canals; fences & fencerows; firebreaks; golf courses; gravel yards; habitat restoration & management areas; highways & roadsides (including aprons, medians, guardrails & right of ways); industrial plant sites; industrial areas; lumber yards; mulched areas; natural areas; paths and trails; parking areas; parks; paved areas; petroleum & other tank farms; pumping installations; pipeline, power, telephone & utility rights-of-way; power stations; preplant to turf & ornamental plants; railroad rights-of way; recreation areas; refineries; resorts; schools; sidewalks; sports areas; storage areas; substations; tennis courts; uncropped farmstead areas; uncultivated non-agricultural areas; vacant lots; walkways; wastelands; & wildlife habitat areas

When applied as directed for "NON-CROP USES", under conditions described, this product may be used to control the listed weeds in terrestrial non-crop sites within these areas: Habitat Restoration & Management Area, Pipeline, Power, Telephone & Utility Rights-of-Way and Pumping Installations.

This product is a non-selective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks. Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. This product has no herbicide activity in the soil and will not wash or leach to affect nearby vegetation. Any ornamental species may be planted in treated areas 7 days or more after application. For most effective results, delay mowing, clipping, planting or sodding of treated areas for at least 7 days after application. This allows time for this product to move within the plant.

For specific rates of application and instructions for control of particular annual weeds, perennial weeds, woody brush and trees, see the "WEEDS CONTROLLED" section of this label. These applications may be made to large affected areas or as spot treatments. For general use in small areas, see alternative instructions below under "Small Area Treatment With Hand-held Sprayers".

Unless the "Agriculture Use Requirements" on this label are observed, the following restrictions apply:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS. IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.

Depending on the type of non-crop application, this product may be applied with boom equipment, high-volume spray equipment and hand-held sprayers as described in the respective portions of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label. Additionally, the product may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any non-crop site specified on this label. See the "Selective Equipment" part of "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Small Area Treatment With Hand-held Sprayers

Add 2.25 to 4.5 fluid ounces of this product plus 0.5 to 1 fluid ounce of nonionic surfactant to 1 gallon of clean water. Use the low rate for many grasses and annual weeds. Use the higher specified rate for control of perennials and brush. Use pump-up sprayer, backpack sprayer or other sprayer suitable for small areas. Adjust equipment to deliver a coarse spray pattern. USE OF HOSE-END SPRAYERS OR SPRINKLER-TYPE DEVICES MAY NOT BE USED.

TANK MIXTURES FOR NON-CROP SITES

When applied as a tank mixture, this product provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide preemergence control of the weeds listed in the individual product labels.

This product PLUS Diuron
This product PLUS Krovar® I
This product PLUS Princep®, Caliber®90, Simazine 4L, 80W or 90DF
This product PLUS Surflan®75W, Surflan AS
This product PLUS Ronstar®50WP
This product PLUS Spyder or Spyder Extra
This product PLUS ProClipse
This product PLUS Polaris AC Complete

When tank mixing with residual herbicides, add a nonionic surfactant at 0.5 to 1 percent by volume of spray solution. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label before preparing these tank mixtures.

Read and carefully observe the label claims, precautionary statements, specified use rate and all other information on the labels of all products used in these tank mixtures.

Use according to the most restrictive label directions for each product in the mixture.

CONTROL OF EMERGED WEEDS

Note: For backpack sprayer and handgun applications, see the "HAND-HELD AND HIGH VOLUME EQUIPMENT" section for specified rates.

Annual Weeds

Apply 1.5 pints per acre of this product in these tank mixtures when weeds are less than 6 inches tall and 2.25 pints per acre when weeds are more than 6 inches tall.

Perennial Weeds

For partial control of perennial weeds using these tank mixtures, apply 1.5 to 7.5 pints per acre of this product. Follow the recommendations in the "WEEDS CONTROLLED" section of this label for stage of growth and rate of application for specific perennial weeds.

PREEMERGENCE WEED CONTROL

For preemergence weed control, refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.

Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

BROADCAST APPLICATION FOR WEED CONTROL IN CHRISTMAS TREE PLANTATIONS

NOTE: IF THIS PRODUCT IS IMPROPERLY APPLIED, IT HAS THE POTENTIAL TO CAUSE SEVERE INJURY TO CHRISTMAS TREES. FOLLOW ALL LABELED DIRECTIONS.

This product may be applied as a broadcast spray over established Christmas trees. To prevent drift onto nearby desirable crops or vegetation, ensure that adequate buffers are maintained.

The following Christmas tree species are approved for this application:

- Douglas Fir (*Pseudotsuga menziesii*)
- Fir species (*Abies* spp.)
- Spruce species (*Picea* spp.)

Do not apply this product until trees have completed at least a full growing season since planting or transplanting. Do not apply within 1 full year prior to tree harvest.

In the fall, applications may only be made after the formation of final conifer resting buds. Final resting buds must be in the dormant stage and fully hardened. If applications are made at any other time, unacceptable Christmas tree injury may occur.

Avoid spray pattern overlap, as injury may result.

Apply 24 fluid ounces of this product per acre in 5 to 30 gallons of water per acre.

NOTE: ADDING SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO THIS PRODUCT MAY RESULT IN SEVERE CHRISTMAS TREE INJURY.

In some areas, this product may be used at rates from 24 to 48 fluid ounces per acre. Consult your local Nufarm representative for specific instructions if you require rates that exceed 24 fluid ounces per acre.

Do not use drift control additives as they may increase Christmas tree injury. Do not use other herbicides in a tank mix with this product as Christmas trees could be severely injured.

SILVICULTURAL SITES AND RIGHTS-OF-WAY

NOTE: DO NOT USE AS AN OVER-THE-TOP BROADCAST SPRAY IN SILVICULTURAL NURSERIES.

When applied as directed for "NON-CROP USES" under conditions described this product controls undesirable vegetation listed on this label. This product also suppresses or controls undesirable vegetation listed on this label when applied at specified rates for release of established coniferous species listed on this label.

For specific rates of application and instructions for control of various brush, annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. For specific rates of application for release of listed coniferous species, see the "CONIFER RELEASE" part of this section of the label.

Where repeat applications are necessary, do not exceed 8 quarts of this product per acre per year.

Aerial Application

This product may be applied using aerial spray equipment for silvicultural site preparation, conifer release and rights-of-way treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING AND APPLICATION INSTRUCTIONS APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on how to apply this product by air. DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

SITE PREPARATION

Following preplant applications of this product, any silvicultural species may be planted.

POST DIRECTED SPRAY

In established silvicultural sites, use as a spray on the foliage of undesirable vegetation. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

CONIFER RELEASE

For release, apply at the end of the first growing season, except in California. Vegetation of target weeds or trees should not be disturbed prior to treatment or until visual symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late Fall. **Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of this label.

For release of the following conifer species:

Douglas Fir <i>Pseudotsuga menziesii</i>	Fir <i>Abies</i> spp.	Hemlock <i>Tsuga</i> spp.	Pines* <i>Pinus</i> spp.	Spruce <i>Picea</i> spp.
--	---------------------------------	-------------------------------------	------------------------------------	------------------------------------

*Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 2.25 to 3 pints of this product per acre except in Washington and Oregon, west of the crest of the Cascade Mountains. For Spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For Fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1.5 to 2.25 pints of this product per acre before any major leaf drop of deciduous species. Add 10 fluid ounces nonionic surfactant per 2 pints of this product. In Maine, up to 4.5 pints per acre may be used for the control of difficult weeds.

Note for Douglas fir release: Ensure that surfactant has been adequately tested for Douglas fir safety and follow manufacturer's specifications for rate of application.

For release of Western hemlock, apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly Pine <i>Pinus taeda</i>	Eastern white pine <i>Pinus strobus</i>	Slash pine <i>Pinus elliottii</i>
--	---	---

Late Season Application - Apply 2-1/4 to 3 pints of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Nufarm does not recommend the use of a crop oil concentrate or MSO (methylated seed oil) based surfactant for use in southern conifer species release with this product. The addition of a tested and approved southern conifer release surfactant is recommended. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will create the potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later applications. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants.

Applications made according to label directions will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash <i>Fraxinus</i> spp.	Hawthorn <i>Crataegus</i> spp.	Oak, Post <i>Quercus stellata</i>	Poplar, yellow <i>Liriodendron tulipifera</i>	Sumac, Smooth <i>Rhus glabra</i>
Cherry, Black <i>Prunus serotina</i>	Locust, Black <i>Robinia pseudoacacia</i>	Oak, Southern Red <i>Quercus falcata</i>	Sassafras <i>Sassafras albidum</i>	Sumac, Winged <i>Rhus copallina</i>
Cherry, Pin <i>Prunus pensylvanica</i>	Maple, Red <i>Acer rubra</i>	Oak, White <i>Quercus alba</i>	Sourwood <i>Oxydendrum arboreum</i>	Sweetgum <i>Liquidambar styraciflua</i>
Elm <i>Ulmus</i> spp.	Oak, Black <i>Quercus velutina</i>	Persimmon <i>Diospyros</i> spp.	Sumac, Poison <i>Rhus vernix</i>	

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

THIS PRODUCT PLUS SPYDER TANK MIXTURES FOR CONIFER RELEASE FROM HERBACEOUS WEEDS

To release Loblolly pines, Slash, Red pine and Virginia pine from herbaceous weeds, tank mixtures of this product with Spyder will provide control of annual weeds listed in the "WEEDS CONTROLLED" section of this and the Spyder label, and partial control of the perennial weeds listed below .

Apply 12 to 18 fluid ounces of this product plus 2 to 4 fluid ounces of Spyder in 10 to 30 gallons of spray solution per acre. Nufarm does not recommend the use of a crop oil concentrate or MSO (methylated seed oil) based surfactant for use in southern conifer species release with this product. The addition of a tested and approved southern conifer release surfactant is recommended. Make application to actively growing weeds as a broadcast spray over the top of the young Loblolly pine, Red pine, Slash pine and Virginia pine.

This tank mixture may be applied using aerial equipment. When applying by air, use the specified rate in 5 to 15 gallons of spray solution per acre. This product plus Spyder tank mixtures may not be applied by air in California.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use the lower rates of both products.

Use the higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass <i>Paspalum notatum</i>	Dock, curly <i>Rumex crispus</i>	Fescues, tall <i>Festuca arundinacea</i>	Poorjoe* <i>Diodia teres</i>	Vaseygrass <i>Paspalum urvillei</i>
Broomsedge <i>Andropogon virginicus</i>	Dogfennel <i>Eupatorium capilliflorum</i>	Johnsongrass* <i>Sorghum halepense</i>	Trumpet creeper** <i>Campsis radicans</i>	Vervain, blue <i>Verbena hastata</i>

*Control at the higher rates

**Suppression at the higher rates only.

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water, insects or disease, or are in an active growth stage.

Read and observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Note To User: This product must not be used in areas where adverse impact on federally designated endangered/threatened plant or aquatic species is likely. Prior to making applications, the user of this product must determine that no such species are located in or immediately adjacent to the area to be treated.

WETLAND SITES

This product may be used in and around water (aquatic areas) and wetlands found in forestry and in power, telephone and pipeline rights-of-way sites including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

Note: Do not apply this product directly to water within 0.5 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 0.5 mile of an active potable water intake in a standing body of water such as a lake, pond or reservoir. To make aquatic applications around and within 0.5 mile of active potable water intakes, the water intake must be turned off for a minimum period

of 48 hours after application. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist. The maximum application rate of 3.75 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any specified rate may be applied:

- Stream crossings in utility right-of-way.
- Where applications will result in less than 20 percent of the total water area being treated.

WILDLIFE HABITAT RESTORATION AND MANAGEMENT AREAS

This product is for the restoration and/or maintenance of native habitat and in wildlife management areas.

Habitat Restoration and Maintenance

When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

Wildlife Food Plots

This product may be used as site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to re-infest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

WIPER APPLICATIONS

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Wiper applications can be used to control or suppress annual and perennial weeds listed on this label. In heavy weed stands, a double application in opposite directions may improve results. See the "WEEDS CONTROLLED" section in this label for specified timing, growth stage and other instructions for achieving optimum results.

CUT STUMP APPLICATION

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. **Apply a 50 to 100 percent solution of this product to the freshly-cut surface immediately after cutting.** Delay in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will control, partially control or suppress many types of woody brush and tree species, some of which are listed below:

Alder <i>Alnus</i> spp.	Eucalyptus <i>Eucalyptus</i> spp.	Maple* <i>Acer</i> spp.	Reed, Giant <i>Arundo donax</i>	Sycamore <i>Platanus occidentalis</i>
Coyote Brush* <i>Baccharis consanguinea</i>	Hickory* <i>Carya</i> spp.	Oak <i>Quercus</i> spp.	Salt cedar <i>Tamarix</i> spp.	
Dogwood* <i>Cornus</i> spp.	Madrone <i>Arbutus menziesii</i>	Poplar <i>Populus</i> spp.	Sweet gum <i>Liquidambar styraciflua</i>	

*This product is not approved for this use on this species in the state of California.

INJECTION AND FRILL APPLICATIONS

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 1 ml of this product per 2 to 3 inches of trunk diameter. This is best achieved by applying 25 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying dilute material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion.

Control		Suppression	
Oak	<i>Quercus</i> spp.	Black Gum*	<i>Nyssa sylvatica</i>
Poplar	<i>Populus</i> spp.	Dogwood	<i>Cornus</i> spp.
Sweetgum	<i>Liquidambar styraciflua</i>	Hickory	<i>Carya</i> spp.
Sycamore	<i>Platanus occidentalis</i>	Maple, Red	<i>Acer rubrum</i>

*This product is not approved for this use on this species in the state of California.

**INJECTION METHOD FOR CONTROL OF JAPANESE KNOTWEED (*Polygonum cuspidatum*)
& GIANT KNOTWEED (*Polygonum polystachyum*)**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the AquaNeat Herbicide label booklet must be followed.

See the "PRODUCT INFORMATION" and "MIXING AND APPLICATION INSTRUCTIONS" sections of this product's label booklet for essential product performance information.

This product may be used for control of Japanese knotweed and giant knotweed using individual stem treatment. Individual knotweed stems may be treated by injecting up to 5 ml of this product, undiluted directly into the hollow stem just below a node. A hole suitable for injecting the herbicide should be made through both sides of the stem using an awl or other convenient pointed tool about 6 inches above the ground, just below a node. (Nodes are circular thickenings or scars surrounding the stem where leaves are or were previously attached.) The herbicide is then injected into this hole. Each stem of the knotweed plant must be treated.

This product can be injected using any injection device capable of delivering a 5 ml dose. For convenience and accuracy, a hand-operated injection device designed to deliver repeated pre-measured doses from a supply reservoir is recommended.

Commercially available dose measuring equipment may be adapted for this purpose. Calibrate the device to deliver a dose of 5 ml per injection cycle. A sharpened hollow probe for puncturing the stem and delivery of the herbicide can also be integrated into the delivery system.

Restriction: Do not apply more than 7.5 quarts of this product per acre. At 5 ml per stem, 7.5 quarts is sufficient to treat a maximum of 1,420 stems per acre.

**RELEASE OF BERMUDAGRASS OR BAHIAGRASS ON NONCROP SITES
RELEASE OF DORMANT BERMUDAGRASS AND BAHIAGRASS**

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Make applications to dormant bermudagrass or bahiagrass.

For best results on winter annuals, treat when weeds are in an early growth stage (below 6 inches in height) after most have germinated.

For best results on tall fescue, treat when fescue is in or beyond the 4- to 6-leaf stage.

WEEDS CONTROLLED

Rate for control or suppression of winter annuals and tall fescue are listed below.

Apply the specified rates of this product in 10 to 25 gallons of water per acre, plus 2 quarts nonionic surfactant per 100 gallons of total spray volume.

WEEDS CONTROLLED OR SUPPRESSED*

**NOTE: C = Control
S = Suppression**

WEED SPECIES	AQUANEAT AQUATIC HERBICIDE (FLUID OZ/ACRE)					
	6	9	12	18	24	48
Barley, little <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annual</i>	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>	.	S	S	C	C	C
Clover, large hop <i>Trifolium campestre</i>	.	S	S	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	C	C	C	C	C
Fescue, tall <i>Festuca arundinacea</i>	S	S
Geranium, Carolina <i>Geranium carolinianum</i>	.	.	S	S	C	C
Henbit <i>Lamium amplexicaule</i>	.	S	C	C	C	C
Ryegrass, Italian <i>Lolium multiflorum</i>	.	.	S	C	C	C
Vetch, common <i>Vicia sativa</i>	.	.	S	C	C	C

*These rates apply only to sites where an established competitive turf is present.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

NOTE: USE ONLY ON SITES WHERE BAHIAGRASS OR BERMUDAGRASS ARE DESIRED FOR GROUND COVER AND SOME TEMPORARY INJURY OR YELLOWING OF THE GRASSES CAN BE TOLERATED.

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "WEEDS CONTROLLED" section in this label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed in this label, use 3/4 to 2-1/4 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre, plus 2 quarts of a nonionic surfactant per 100 gallons of total spray volume. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as size of plants increases or as they approach flower or seedhead formation.

Use the higher rate for partial control or longer-term suppression of the following perennial species. Use lower rates for shorter-term suppression of growth.

Bahiagrass Johnsongrass**
Dallisgrass Trumpetcreeper*
Fescue (tall) Vaseygrass

*Suppression at the higher rate only.

**Johnsongrass is controlled at the higher rate.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When applied as directed in the "NONCROP SITES" section in this label, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full green-up of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 5 fluid ounces per acre of this product, plus 2 quarts of an approved nonionic surfactant per 100 gallons of total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued vegetative growth suppression, sequential applications must be made prior to seedhead emergence.

Apply no more than 2 sequential applications per year. As a first sequential application, apply 3 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 3 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

ANNUAL GRASS GROWTH SUPPRESSION

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 ounces of this product in 10 to 40 gallons of spray solution per acre. Mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

AQUATIC SITES

When applied as directed and under the conditions described in the "WEEDS CONTROLLED" section in this label, this product will control or partially control the labeled weeds growing in aquatic sites.

Aquatic Sites - This product may be applied to emerged weeds in all bodies of fresh and brackish water which may be flowing, non-flowing or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas, and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

This product does not control plants which are completely submerged or have a majority of their foliage under water.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.

NOTE: Do not apply this product directly to water within 1/2 mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made **ONLY** in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds. Floating Mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not re-treat within 24 hours following the initial treatment.

Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 7-1/2 pints per acre must not be exceeded in any single broadcast application that is being made over water.

When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store below 32°F or above 100°F. Store in original container in a well-ventilated area separately from fertilizer, feed, and food stuffs. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL [HANDLING]:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container disposal [handling] instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV072413)

AquaNeat is a registered trademark of Nufarm, Inc.